

## United States Patent [19]

Akhavan-Tafti

[11] Patent Number:

6,001,614

[45] Date of Patent:

\*Dec. 14, 1999

# [54] METHODS OF SYNTHESIZING LABELED POLYNUCLEOTIDES BY LIGATION OF MULTIPLE OLIGOMERS

[75] Inventor: Hashem Akhavan-Tafti, Howell, Mich.

[73] Assignee: Lumigen, Inc., Southfield, Mich.

[\*] Notice: This patent is subject to a terminal dis-

claimer.

[21] Appl. No.: 09/241,353

[22] Filed: Feb. 2, 1999

#### Related U.S. Application Data

[63]	Continuation	of application	No.	09/121,887, Jul.	24,	1998.
------	--------------	----------------	-----	------------------	-----	-------

[51] Int. Cl.<sup>6</sup> ...... C12P 19/34; C12Q 1/68; C07H 21/04

#### [56] References Cited

#### U.S. PATENT DOCUMENTS

		Brennan
5,663,062	9/1997	Sorge 435/91.1
		Schalling 435/91.52
5,750,341		Macevicz 435/6
5,770,367		Southern
5,888,731	3/1999	Yager 435/6

### FOREIGN PATENT DOCUMENTS

88311741 6/1989 European Pat. Off. .

#### OTHER PUBLICATIONS

S. Dubiley, E. Kirilov, Y. Lysov, A. Mirzabekov Nucl. Acids Res., 25, 2259–2265 (1997). R. Handley, H. Akhavan–Tafti, A. P. Schaap, J. Clin. Ligand Assay, 20(4) 302–312 (1997). K.D.James, A.D.Ellington, Chemistry & Biology 4, 595-605 (1997).

T.Kaczorowski, W.Szybalski, Gene 179, 189-193 (1996).

L.E.Kotler, D.Zevin-Sonkin, I.A.Sobolev, A.D.Beskin L.E.Ulanovsky, Proc. Natl. Acad. Sci. USA, 90, 4241-4245 (1993).

T.Li, K.C. Nicolaou, Nature, 369, 218-221 (1994).

C.E.Pritchard,E.M.Southern,Nucl.Acids Res., 25, 3403-3407 (1997).

D.Sievers, G. Von Kiedrowski, Nature, 369, 221-224 (1994).

W.Stemmer, A.Crameri, K.D.Ha, T.M.Brennan, H.L. Heyneker, Gene, 164, 49-53 (1995).


D.Y.Wu, R.B. Wallace, Genomics, 4,560-569 (1989).

Primary Examiner—Carla J. Myers
Assistant Examiner—Diana Johannsen
Attorney, Agent, or Firm—Richard S. Handley

#### [7] ABSTRACT

Methods of synthesizing polynucleotides are disclosed involving the simultaneous ligation of a set of oligomer 5'-phosphates onto a template-bound primer. The set of these oligomers can be preselected to contain oligomers which are complementary to the template strand or the oligomers can be supplied as a library and allowed to self select. The synthesis by ligation can proceed unidirectionally or bidirectionally from the primer and can be used to synthesize both strands simultaneously by the use of two primers. The ligation is preferably performed with a ligase enzyme. The methods of synthesis are useful in a variety of applications, including cloning, amplification, labeling, diagnostic assays, mutation analysis and screening, gene expression monitoring and sequence analysis.

23 Claims, 6 Drawing Sheets



•